

PCT

WORLD INTELLECTUAL PROPERTY ORGANIZATION
International Bureau




INTERNATIONAL APPLICATION PUBLISHED UNDER THE PATENT COOPERATION TREATY (PCT)

(51) International Patent Classification ⁶ : C07D 455/04, 215/22, A61K 31/435		A1	(11) International Publication Number: WO 97/29107
			(43) International Publication Date: 14 August 1997 (14.08.97)
(21) International Application Number: PCT/EP97/00552 (22) International Filing Date: 7 February 1997 (07.02.97) (30) Priority Data: FI96A000019 9 February 1996 (09.02.96) IT (71) Applicant (for all designated States except US): APPLIED RESEARCH SYSTEMS ARS HOLDING N.V. [NL/NL]; John B. Gorsiraweg 14, Curaçao (AN). (72) Inventors; and (75) Inventors/Applicants (for US only): GUARNA, Antonio [IT/IT]; Via Pistoiese, 158, I-50040 Seano (IT). SERIO, Mario [IT/IT]; Via Roma, 41, I-52010 Subbiano (IT). (74) Agent: GERVASI, Gemma; Notarbartolo & Gervasi, Corso di Porta Vittoria, 9, I-20122 Milan (IT).		(81) Designated States: AL, AM, AT, AU, AZ, BA, BB, BG, BR, BY, CA, CH, CN, CU, CZ, DE, DK, EE, ES, FI, GB, GE, HU, IL, IS, JP, KE, KG, KP, KR, KZ, LC, LK, LR, LS, LT, LU, LV, MD, MG, MK, MN, MW, MX, NO, NZ, PL, PT, RO, RU, SD, SE, SG, SI, SK, TJ, TM, TR, TT, UA, UG, US, UZ, VN, YU, ARIPO patent (KE, LS, MW, SD, SZ, UG), Eurasian patent (AM, AZ, BY, KG, KZ, MD, RU, TJ, TM), European patent (AT, BE, CH, DE, DK, ES, FI, FR, GB, GR, IE, IT, LU, MC, NL, PT, SE), OAPI patent (BF, BJ, CF, CG, CI, CM, GA, GN, ML, MR, NE, SN, TD, TG). Published With international search report. Before the expiration of the time limit for amending the claims and to be republished in the event of the receipt of amendments.	

(54) Title: BENZO[C]QUINOLIZINE DERIVATIVES, THEIR PREPARATION AND USE AS 5 α -REDUCTASES INHIBITORS

(57) Abstract

The present invention refers to benzo[c]quinolizine derivatives of general formula (I), their pharmaceutically acceptable salts or esters, processes for their preparation and pharmaceutical compositions containing them.



FOR THE PURPOSES OF INFORMATION ONLY

Codes used to identify States party to the PCT on the front pages of pamphlets publishing international applications under the PCT.


AM	Armenia	GB	United Kingdom	MW	Malawi
AT	Austria	GE	Georgia	MX	Mexico
AU	Australia	GN	Guinea	NE	Niger
BB	Barbados	GR	Greece	NL	Netherlands
BE	Belgium	HU	Hungary	NO	Norway
BF	Burkina Faso	IE	Ireland	NZ	New Zealand
BG	Bulgaria	IT	Italy	PL	Poland
BJ	Benin	JP	Japan	PT	Portugal
BR	Brazil	KE	Kenya	RO	Romania
BY	Belarus	KG	Kyrgyzstan	RU	Russian Federation
CA	Canada	KP	Democratic People's Republic of Korea	SD	Sudan
CF	Central African Republic	KR	Republic of Korea	SE	Sweden
CG	Congo	KZ	Kazakhstan	SG	Singapore
CH	Switzerland	LI	Liechtenstein	SI	Slovenia
CI	Côte d'Ivoire	LK	Sri Lanka	SK	Slovakia
CM	Cameroon	LR	Liberia	SN	Senegal
CN	China	LT	Lithuania	SZ	Swaziland
CS	Czechoslovakia	LU	Luxembourg	TD	Chad
CZ	Czech Republic	LV	Latvia	TG	Togo
DE	Germany	MC	Monaco	TJ	Tajikistan
DK	Denmark	MD	Republic of Moldova	TT	Trinidad and Tobago
EE	Estonia	MG	Madagascar	UA	Ukraine
ES	Spain	ML	Mali	UG	Uganda
FI	Finland	MN	Mongolia	US	United States of America
FR	France	MR	Mauritania	UZ	Uzbekistan
GA	Gabon			VN	Viet Nam

- 1 -

Benzo[c]quinolizine derivatives, their preparation and use as 5 α -reductases inhibitors.

Field of the invention

The present invention refers to benzo[c]quinolizine derivatives of
5 general formula (I)



wherein:

R₁, R₂, R₃, R₄, R₆, same or different, are chosen in the group consisting of: H, C₁₋₈alkyl, C₂₋₈alkenyl, C₂₋₈alkinyl, cycloalkyl, aryl, heterocycle, halogen, CN, azide, NRR', C₁₋₈alkylamino,
10 arylamino, C₁₋₈alkyloxy, aryloxy, COOR, CONRR' wherein R and R', same or different, are chosen in the group consisting of: H, C₁₋₈alkyl, cycloalkyl, aryl, heterocycle, arylC₁₋₈alkyl;

R₅ is chosen in the group consisting of: H, C₁₋₈alkyl, COOR, CN, aryl, heterocycle;

15 X is chosen in the group consisting of: O, C(=O)R, COOR, NO₂, CONR'R wherein R and R' are as above defined;

Q is chosen in the group consisting of: simple bond, C₁₋₈alkyl, C₂₋₈alkenyl, C₂₋₈alkinyl, cycloalkyl, CO, CONR, NR, wherein R is as above defined;

20 W is chosen in the group consisting of: H, C₁₋₈alkyl, C₂₋₈alkenyl, C₂₋₈alkinyl, cycloalkyl, trifluoromethyl, C₁₋₈alkoxy, C₁₋₈alkoxy-C₁₋

- 2 -

galkyl, arylC₁₋₈galkyl, aryl, aryloxy, arylamino, C₁₋₈galkylcarbonyl, arylcarbonyl, halogen, CN, NRR', C₁₋₈galkylamino, heterocycle wherein the groups alkyl, alkenyl, alkynyl, cycloalkyl, aryl, heterocycle, can be substituted;

5 n is an integer comprised between 1 and 4;

the symbol ----- means that the corresponding bonds a, b, c, d e, f, and g can be simple or double bonds;

with the proviso that when b or f are a double bond then the group R₅ is absent;

10 their pharmaceutically acceptable salts or esters, their process of preparation and their use as inhibitors of steroid 5 α -reductases (hereinafter indicated as 5 α -reductases).

State of the art

The enzyme known as steroid 5 α -reductase is a system formed by two
15 iso-enzymes (type I and type II or 5 α R-I and 5 α R-II respectively)) which converts testosterone into dihydrotestosterone, the most powerful androgen circulating in the body.

The type I iso-enzyme (5 α R-I) is mainly present in liver and skin while the type II iso-enzyme (5 α R-II) is mainly present in the
20 prostate tissue and in the male sexual organs and its activity is essential in the fetal developing process for the differentiation of the external sexual organs.

The production of dihydrotestosterone is associated with some pathologies which are widely diffused as for example benign prostate
25 hypertrophy, prostate cancer, baldness and acne in men and hirsutism in women. More particularly iso-enzyme I plays a role in the pathologies regarding the skin while iso-enzyme-II is involved in

- 3 -

prostate pathologies.

In the recent years a lot of international searchers have tried to isolate new compounds capable of inhibiting the 5 α -reductase enzyme in order to treat the above said pathologies, especially, if possible,


5 acting selectively on only one of the two iso-enzymes.

Inhibitors of 5 α -reductase, and also of the iso-enzymes 5 α R-I and 5 α R-II were already described, for example finasteride used with success in the treatment of benign prostate hypertrophy [see for example J.Med.Chem. 36, 4313-15 (1993), J.Med.Chem. 37, 3871-74 (1994)]. It is
10 therefore evident the importance of developing new compounds capable of inhibiting the action of the 5 α -reductase enzyme and in particular capable of acting selectively on 5 α R-I iso-enzyme which, as said, is responsible, of widely diffused pathologies having an high impact as baldness in men and hirsutism in women.

15 Detailed description of the invention

The present invention refers to new compounds capable of inhibiting the 5 α -reductase enzyme, either selectively in respect of 5 α R-I and 5 α R-II or on both the iso-enzymes, useful for the treatment of the pathologies mediated by the enzyme.

20 The products according to the invention have general formula (I)



- 4 -

wherein the substituents R_1 , R_2 , R_3 , R_4 , R_5 , R_6 , X, Q, W, n and the symbol ----- are as above defined.

According to the present invention with group C_{1-8} alkyl, C_{2-8} alkenyl and C_{2-8} alkinyl are indicated linear or branched alkyl radicals as for
5 example: methyl, ethyl, propyl, isopropyl, butyl, pentyl, hexyl, heptyl, octyl, ethylene, propene, butene, isobutene, acetylene, propine, butine ecc.

With cycloalkyl are indicated: cyclopropane, cyclobutane, cyclopentane, cyclohexane, cycloheptane, cyclooctane, norbornane,
10 canphane, adamantane.

With aryl are indicated: phenyl and naphthyl.

Heterocycle means in particular: saturated or aromatic heterocycles containing one or more N atoms, more particularly: piridine, imidazole, pyrrole, indole, triazoles, pyrrolidine, piperidine.

15 Halogen means: fluorine, chlorine, bromine, iodine.

The substituents of the above said group W are preferably: halogen, OR, phenyl, NRR' , CN, COOR, CONRR', C_{1-8} alkyl (wherein R and R' are as above defined).

In particular, according to the present invention compounds of formula

20 (I) are preferred wherein:

R_5 = H, heterocycle

X = O

Q = simple bond, CO, CONR, NR (wherein R is as above defined)

W = H, F, Cl, Br, Me, t-butyl, C_{1-8} alkoxy, 2,5-dimethylhexyl,
25 trifluoromethyl, 2,5-(di-trifluoromethyl)-phenyl, 4-methoxy-phenyl, 4-fluoro-phenyl, phenyl, phenyl- C_{1-8} alkyl, C_{1-8} alkylcarbonyl, phenylcarbonyl.

- 5 -

$n = 1$ and 2

$R_1, R_2, R_3, R_4, R_6 = H, Me, CN, \text{phenyl}, COOR, CONRR'$ (wherein R and R' are as above defined).

Among the pharmaceutically acceptable esters and salts according to
5 the present invention the following can be mentioned: hydrochloride, sulphate, citrate, formate, phosphate.

Preferred compounds according to the present invention are:


- 1,2,4,4a,5,6 hexahydro-(11H)-benzo[c]quinolizine-3-one;
- 8-chloro-1,2,4,4a,5,6 hexahydro-(11H)-benzo[c]quinolizine-3-one;
- 10 1,2,4,4a,5,6 hexahydro-8-methyl-(11H)-benzo[c]quinolizine-3-one;
- 1,2,4,4a,5,6 hexahydro-4-methyl-(11H)-benzo[c]quinolizine-3-one;
- 1,2,4,4a,5,6 hexahydro-1-methyl-(11H)-benzo[c]quinolizine-3-one;
- 1,2,5,6 tetrahydro-(11H)-benzo[c]quinolizine-3-one;
- 8-chloro-1,2,5,6 tetrahydro-(11H)-benzo[c]quinolizine-3-one;
- 15 8-methyl-1,2,5,6 tetrahydro-(11H)-benzo[c]quinolizine-3-one;
- 4-methyl-1,2,5,6 tetrahydro-(11H)-benzo[c]quinolizine-3-one;
- 1-methyl-1,2,5,6 tetrahydro-(11H)-benzo[c]quinolizine-3-one;
- 4,4a,5,6 tetrahydro-(11H)-benzo[c]quinolizine-3-one;
- 5,6-dihydro-(11H)-benzo[c]quinolizine-3-one;
- 20 8-chloro-4,4a,5,6 tetrahydro-(11H)-benzo[c]quinolizine-3-one;
- 8-chloro-1-methyl-4,4a,5,6-tetrahydro-(11H)-benzo[c]quinolizine-3-one;
- 8-methyl-4,4a,5,6 tetrahydro-(11H)-benzo[c]quinolizine-3-one;
- 4-methyl-4,4a,5,6 tetrahydro-(11H)-benzo[c]quinolizine-3-one (cis) and (trans);
- 25 8-chloro-4-methyl-1,2,5,6 tetrahydro-(11H)-benzo[c]quinolizine-3-one;
- 4,8-dimethyl-1,2,5,6 tetrahydro-(11H)-benzo[c]quinolizine-3-one;
- 4,8-dimethyl-4,4a,5,6 tetrahydro-(11H)-benzo[c]quinolizine-3-one (cis)

- 6 -

and (trans);

8-chloro-4-methyl-4,4a,5,6-tetrahydro-(11H)-benzo[c]quinolizine-3-one
(cis) and (trans).

The compounds according to the present invention can be prepared for
5 example starting from compounds of formula 2



(2)

wherein R_3 , R_4 , W, Q and n are as above defined, following the
reaction Scheme reported hereinafter.

The compounds 2 are commercially available or can be prepared according
to known techniques.

- 10 As it can be seen from the Scheme the preparation of the compounds
according to the invention involves the protection of the amide-group
in compound 2 by the protecting group Z, for example tert-
butoxycarbonyl (t-Boc), to give compound 3; compound 3 is reduced to
compound 4, for example (when R_5 is H) with sodium borohydride in
15 ethanol (pH 3), which is reacted with a silylether 6, produced "in
situ" starting from vinyl-ketones 5 (wherein R_1 , R_2 and R_6 are as
above defined) with a silylating agent as
trimethylsilyltrifluoromethanesulphonic anhydride (TMSOTf) and
thereafter hydrolyzed, for example in sodium hydrogencarbonate, to
20 give the compounds of formula (I) wherein $X = O$. The possible
introduction of the double bonds and the transformation of the group X

- 7 -

in one of the other groups mentioned above can be easily performed according to known techniques starting from the corresponding compound of formula (I) obtained as indicated. For example the introduction of the double bonds in position a or b, can be performed by reaction of 5 dichlorodicyanoquinone (DDQ) with the corresponding silylenolethers or by oxidation with mercuric acetate of the saturated corresponding compound obtained as described above. The transformation of group X can be performed via the corresponding enoltriflates and their carbonylation in the presence of palladium diacetate, 10 triphenylphosphine and the suitable nucleophilic reagent (alcohol, amine, nitro-group).

Example 1

Preparation of N-(t-butoxycarbonyl)-3,4-dihydroquinolin-2(1H)-one

[compound 3 wherein $(QW)_n = H$, $R_3 = R_4 = H$]

15 5 g (34 mmol) of 3,4-dihydroquinolin-2(1H)-one [compound 2 wherein $(QW)_2 = H$, $R_3 = R_4 = H$] and 111 ml of CH_2Cl_2 are charged, under nitrogen, in a 250 ml round bottom flask, equipped with magnetic stirrer.

To the above said mixture 4.7 ml (34 mmol) of triethylamine 20 (distilled on KOH), 8.9 g (40.8 mmol) of di-butyl dicarbonate and 1 g (8.2 mmol) of 4-dimethylaminopyridine are added. The mixture is stirred under reflux for 5 h, then for one night at room temperature and thereafter the solvent is removed and 200 ml of water are added. The aqueous phase is extracted with diethylether and the organic phase 25 is neutralized with an aqueous solution of $KHSO_4$ (1 M). The organic phase is washed with a saturated solution of NaCl and dried on Na_2SO_4 . After filtration and removal of the solvent 8.23 g of the desired

product are obtained (white crystals). M.p.: 68 - 69°C. Yield: 98%.

Example 2

Preparation of N-(t-butoxycarbonyl)-2-ethoxy-1,2,3,4-tetrahydroquinoline [compound 4 wherein $(QW)_n = H$, $R_3 = R_4 = R_5 = H$].

5 4.35 g (17.6 mmol) of the compound obtained from example 1 and 136 ml of absolute ethanol are charged in a 500 ml round bottom flask equipped with magnetic stirrer.

The solution is cooled at -25°C and 2.66 g (70.4 mmol) of $NaBH_4$ (subdivided in 6 portions) are added to the mixture in 1 h. After 4 h
10 a solution of HCl 2N in absolute ethanol is added to the mixture, up to pH 3, and the mixture is stirred at 0°C for 1.5 h. 100 ml of water are added, the aqueous phase is extracted with methylene chloride, the organic phase is washed with a saturated solution of $NaHCO_3$ and a saturated solution of NaCl and the mixture is dried on Na_2SO_4 . After
15 filtration the solvent is removed and 4.74 g of the expected product are obtained (dense yellow liquid); yield 96%.

Operating as above said other compounds 4 wherein the substituents can not be reduced by $NaBH_4$ are obtained; if substituents which could be reduced by $NaBH_4$ are present these must be previously protected.

20 Example 3

Preparation of 1,2,4,4a,5,6-hexahydro-(11H)-benzo[c]quinolizin-3-one [compound of formula (I) wherein $X = O$; $(QW)_n = H$; $R_1 = R_2 = R_3 = R_4 = R_5 = R_6 = H$; a,b,c,e,f,g = simple bond]

70 μ l (0.86 mmol) of 3-buten-2-one [compound of formula 5 wherein $R_1 = R_2 = R_6 = H$] and 2 ml of anhydrous CH_2Cl_2 are charged, at 0°C under
25 argon in a two-necked round bottom flask equipped with magnetic stirrer and dropping funnel.

- 9 -

170 μ l (1.22 mmoles) of triethylamine (distilled on KOH) and 209 μ l (1.08 mmoles) of trimethylsilyltrifluoromethanesulphonate (TMDOTf) (drop by drop) are added to the mixture. In this conditions 2-(trimethylsilyloxy)-1,3-butadiene [compound 6 wherein $R_1 = R_2 = R_6 =$
5 H] is formed "in situ". The mixture is stirred for 45 minutes and thereafter a solution of 100 mg (0.36 mmoles) of the product from Example 2 in 2 ml of anhydrous CH_2Cl_2 is added therein, drop by drop, together with 69 μ l (0.36 mmoles) of TMSOTf. The mixture is brought to room temperature and after 30 minutes 4 ml of a saturated solution of
10 NaHCO_3 are added and the mixture is stirred vigorously for 36 h.

4 ml of water are added to the mixture and the aqueous phase is extracted with methylene chloride, the organic phase is washed with a saturated solution of NaHCO_3 , water, a saturated solution of NaCl and is dried on Na_2SO_4 . After filtration the solvent is removed and 59 mg
15 of crude product are obtained. The product is purified by flash chromatography on silica gel column (FCC) eluting with methylene chloride and triethylamine 1%. 18 mg of the wanted product are obtained (crystals). M.p.: 53 - 54°C. Yield 25%.

Using various vinyl-ketones 5, or using directly the various
20 silylenolethers 6 (when available), it is possible to prepare the corresponding derivatives of formula (I).

In particular when 1-methoxy-3-(trimethylsilyloxy)-1,3-butadiene (compound 6 wherein $R_1 = \text{MeO}$, $R_2 = \text{H}$, $R_6 = \text{H}$) was used, 4,4a,5,6-tetrahydro-(11H)-benzo[c]quinolizin-3-one (compound I wherein $X = \text{O}$,
25 $(\text{QW})_n = \text{H}$, $R_1 = R_2 = R_3 = R_4 = R_5 = R_6 = \text{H}$, a = double bond; b,c,e,f,g = single bond] was directly obtained according to the synthesis described in the following Example 4.

- 10 -

Example 4

Preparation of 4,4a,5,6-tetrahydro-(11H)-benzo[c]quinolizin-3-one
[compound I wherein $X = O$, $(QW)_n = H$; $R_1 = R_2 = R_3 = R_4 = R_5 = R_6 = H$;
 $a = \text{double bond}$; $b, c, e, f, g = \text{single bond}$].

- 5 To a stirred solution of compound 4 [$(QW)_n = H$, $R_3 = R_4 = H$] (4 g.
14.42 mmol) of the example 3, in 75 ml of anhydrous CH_2Cl_2 under argon
at $-10^\circ C$ is added, dropwise in 7 min, 28.84 ml of a 1M solution of
 $TiCl_4$ in CH_2Cl_2 maintaining the temperature below $-5^\circ C$. Then 1-
methoxy-3-(trimethylsilyloxy)-1,3-butadiene (compound 6, $R_1 = MeO$, R_2
10 $= H$, $R_6 = H$) (3.29 ml, 17.3 mmol) is added by syringe at $0^\circ C$, and the
reaction was left aside at room temperature for 1 h. The reaction
mixture is added, cautiously, with 100 ml of $NaHCO_3$ saturated solution,
and then stirred for 30 min. The organic layer is separated, washed
with water, filtered on Celite and dried over Na_2SO_4 . After removal of
15 the solvent the crude product is purified by flash column
chromatography (eluant light-petroleum ether/ethyl acetate 1:4)
affording 0.72 g (25% yield) of the expected product (white crystals,
m.p.: $135-137^\circ C$).

Example 5

- 20 a) Preparation of 4-methyl-4,4a,5,6-tetrahydro-(11H)-
benzo[c]quinolizine-3-one [compound of formula (I) wherein $X = O$;
 $(QW)_n = H$; $R_1 = R_3 = R_4 = R_5 = R_6 = H$; $R_2 = Me$; $a = \text{double bond}$;
 $b, c, e, f, g = \text{single bonds}$], 4-methyl-1,2,5,6-tetrahydro-(11H)-
benzo[c]quinolizine-3-one [compound of formula (I) wherein $X = O$;
25 $(QW)_n = H$; $R_1 = R_3 = R_4 = R_6 = H$; $R_2 = Me$; $b = \text{double bond}$; a, c, e, f, g
 $= \text{single bonds}$] and 4-methyl-5,6-dihydro-(11H)-benzo[c]quinolizine-3-
one [compound of formula (I) wherein $X = O$; $(QW)_2 = H$; $R_1 = R_3 = R_4 =$

- 11 -

$R_6 = H$; $R_2 = Me$; a,b = double bonds; c,e,f,g = single bonds].

- 1 g (4.64 mmol) of 4-methyl-1,2,4,4a,5,6-hexahydro-(11H)-benzo[c]quinolizine-3-one [compound of formula (I), wherein $X = O$; (QW)_n = H; $R_1 = R_3 = R_4 = R_5 = R_6 = H$; $R_2 = Me$; a,b,c,e,f,g = single bonds, obtained according to example 3 by reaction of compound 4 (wherein (QW)_n = H; $R_3 = R_4 = R_5 = H$) of example 2 and ethylvinylketone (compound 5 wherein $R_1 = R_6 = H$; $R_2 = Me$] and 120 ml of 5% solution (v/v) of glacial acetic acid in water are charged under nitrogen in a two-necked round bottom flask, equipped with magnetic stirrer, refrigerator and dropping funnel. Under vigorous stirring, 7.27 g (18.56 mmol) of tetrasodic salt EDTA and 5.92 g (18.56 mmol) of (CH₃CO₂)₂Hg are added and the reaction mixture is heated at 90°C for 2h. After cooling at room temperature the reaction mixture is added with 120 ml of water and extracted with methylene chloride (4x70 ml).
- 15 The separated organic phase is washed with a saturated solution of NaHCO₃, with a saturated solution of NaCl then dried over Na₂SO₄. After removal of the solvent the crude product is purified by flash chromatography on silica gel by elution with ethylacetate/light petroleum ether 2:1 affording:
- 20 83 mg (10%) (gummy solid) of cis-4-methyl-4,4a,5,6-tetrahydro-(11H)-benzo[c]quinolizine-3-one [compound of formula (I) wherein $X = O$; (QW)_n = H; $R_1 = R_3 = R_4 = R_5 = R_6 = H$; $R_2 = Me$; a = double bond; b,c,e,f,g = single bonds]
- 350 mg (40%) (crystals, m.p.: 148-150°C) of 4 methyl-1,2,5,6-tetrahydro-(11H)-benzo[c]quinolizine-3-one [compound of formula (I) wherein $X = O$; (QW)_n = H; $R_1 = R_3 = R_4 = R_6 = H$; $R_2 = Me$; b = double bond; a,c,e,f,g = single bonds] and
- 25

- 12 -

107 mg (12%) (gummy solid) of 4-methyl-5,6-dihydro-(11H)-benzo[c]quinolizine-3-one [compound of formula (I) wherein $X = O$; $(QW)_n = H$; $R_1 = R_3 = R_4 = R_6 = H$; $R_2 = Me$; a,b=double bonds; c,e,f,g = single bonds].

5 Activity Test

The inhibition potency of the prepared compounds in respect of the iso-enzymes 1 and 2 of 5 α -reductase was determined using tissue samples (for example prostate human tissue) or human cellular systems (for example DU 145 cells) expressing iso-enzymes 2 and 1
10 respectively.

The samples are incubated in the presence of testosterone labelled with tritium and thereafter the quantity of labelled dihydrotestosterone formed in the absence and in the presence of the inhibitor is measured.

- 15 The compounds showed high inhibiting power of 5 α -reductase enzyme (in particular of iso-enzyme 1) with an inhibition higher than 50% at the concentration of 10 - 100 nM.


For the therapeutical administration the compounds according to the invention are prepared in the form of pharmaceutical compositions
20 containing the active principle and the organic or inorganic excipients suitable for the oral, parenteral or topic administration of the compositions. The pharmaceutical compositions can therefore be in the solid form (dragees, suppositories, creams, ointments), liquid form (solutions, suspensions, emulsions) and can possibly contain the
25 stabilizers, conservatives, humectants, emulsifier, buffers or salts used for equilibrating the osmotic pressure which are commonly used in the art.

- 13 -

Generally the administration of the compounds is performed according to the modalities and quantities observed for the known agents used for the same purposes and taking into consideration the age and conditions of the patients.

Claims

- 1 1. Benzo[c]-quinolizine compounds of formula (I)



- 2 wherein:
- 3 R_1 , R_2 , R_3 , R_4 , R_6 , same or different, are chosen in the group
- 4 consisting of: H, C_{1-8} alkyl, C_{2-8} alkenyl, C_{2-8} alkinyl, cycloalkyl,
- 5 aryl, heterocycle, halogen, CN, azide, NRR' , C_{1-8} alkylamino,
- 6 arylamino, C_{1-8} alkyloxy, aryloxy, COOR, CONRR' wherein R and R', same
- 7 or different, are chosen in the group consisting of: H, C_{1-8} alkyl,
- 8 cycloalkyl, aryl, heterocycle, aryl C_{1-8} alkyl;
- 9 R_5 is chosen in the group consisting of: H, C_{1-8} alkyl, COOR, CN, aryl,
- 10 heterocycle;
- 11 X is chosen in the group consisting of: O, $C(=O)R$, COOR, NO_2 , CONR'R
- 12 wherein R and R' are as above defined;
- 13 Q is chosen in the group consisting of: simple bond, C_{1-8} alkyl, C_{2-8}
- 14 galkenyl, C_{2-8} alkinyl, cycloalkyl, CO, CONR, NR, wherein R is as above
- 15 defined;
- 16 W is chosen in the group consisting of: H, C_{1-8} alkyl, C_{2-8} alkenyl, C_{2-8}
- 17 galkinyl, cycloalkyl, trifluoromethyl, C_{1-8} alkoxy, C_{1-8} alkoxy- C_{1-8}
- 18 galkyl, aryl C_{1-8} alkyl, aryl, aryloxy, arylamino, C_{1-8} alkylcarbonyl,
- 19 arylcarbonyl, halogen, CN, NRR' , C_{1-8} alkylamino, heterocycle wherein
- 20 the groups alkyl, alkenyl, alkinyl, cycloalkyl, aryl, heterocycle, can

- 15 -

21 be substituted;
22 n is an integer comprised between 1 and 4;
23 the symbol ----- means that the corresponding bonds a, b, c, d e, f,
24 and g can be simple or double bonds;
25 with the proviso that when b or f are a double bond then the group R₅
26 is absent;

27 their pharmaceutically acceptable salts or esters.

1 2. Benzo[c]-quinolizine compounds of formula (I)

2 R₅ = H, heterocycle

3 X = O

4 Q = simple bond, CO, CONR, NR (wherein R is as above defined)

5 W = H, F, Cl, Br, Me, t-butyl, C₁₋₈alkoxy, 2,5-dimethylhexyl,

6 trifluoromethyl, 2,5-(di-trifluoromethyl)-phenyl, 4-methoxy-phenyl, 4-

7 fluoro-phenyl, phenyl, phenyl-C₁₋₈alkyl, C₁₋₈alkylcarbonyl,

8 phenylcarbonyl.

9 n = 1 and 2

10 R₁, R₂, R₃, R₄, R₅, R₆ = H, Me, CN, phenyl, COOR, CONRR' (wherein R

11 and R' are as above defined).

1 3. Benzo[c]-quinolizine compounds according to Claim 1 of formula

2 1,2,4,4a,5,6 hexahydro-(11H)-benzo[c]quinolizine-3-one;

3 8-chloro-1,2,4,4a,5,6 hexahydro-(11H)-benzo[c]quinolizine-3-one;

4 1,2,4,4a,5,6 hexahydro-8-methyl-(11H)-benzo[c]quinolizine-3-one;

5 1,2,4,4a,5,6 hexahydro-4-methyl-(11H)-benzo[c]quinolizine-3-one;

6 1,2,4,4a,5,6 hexahydro-1-methyl-(11H)-benzo[c]quinolizine-3-one;

7 1,2,5,6 tetrahydro-(11H)-benzo[c]quinolizine-3-one;

8 8-chloro-1,2,5,6 tetrahydro-(11H)-benzo[c]quinolizine-3-one;

9 8-methyl-1,2,5,6 tetrahydro-(11H)-benzo[c]quinolizine-3-one;


- 16 -

- 10 4-methyl-1,2,5,6 tetrahydro-(11H)-benzo[c]quinolizine-3-one;
- 11 1-methyl-1,2,4,6 tetrahydro-(11H)-benzo[c]quinolizine-3-one;
- 12 4,4a,5,6 tetrahydro-(11H)-benzo[c]quinolizine-3-one;
- 13 5,6-dihydro-(11H)-benzo[c]quinolizine-3-one;
- 14 8-chloro-4,4a,5,6 tetrahydro-(11H)-benzo[c]quinolizine-3-one;
- 15 8-chloro-1-methyl-4,4a,5,6-tetrahydro-(11H)-benzo[c]quinolizine-3-one;
- 16 8-methyl-4,4a,5,6 tetrahydro-(11H)-benzo[c]quinolizine-3-one;
- 17 4-methyl-4,4a,5,6 tetrahydro-(11H)-benzo[c]quinolizine-3-one (cis) and
- 18 (trans);
- 19 8-chloro-4-methyl-1,2,5,6 tetrahydro-(11H)-benzo[c]quinolizine-3-one;
- 20 4,8-dimethyl-1,2,5,6 tetrahydro-(11H)-benzo[c]quinolizine-3-one;
- 21 4,8-dimethyl-4,4a,5,6 tetrahydro-(11H)-benzo[c]quinolizine-3-one (cis)
- 22 and (trans);
- 23 8-chloro-4-methyl-4,4a,5,6-tetrahydro-(11H)-benzo[c]quinolizine-3-one
- 24 (cis) and (trans).

1 4. Process for the preparation of compounds according to Claim 1


2 wherein:

3 the amide-group of a compound of formula (2)



(2)


4 is protected with a protecting group Z to give the compound (3)



(3)

- 17 -

5 the above said compound (3) is reduced to compound (4), for example
 6 with sodium borohydride in ethanol (pH3)



7 and compound (4) is reacted with a silylether (6)



8 prepared "in situ" by reacting a vinyl-ketone (5)




9 (wherein R₁, R₂, R₆ are as above defined) with a silylating agent as
 10 trimethylsilyltrifluorometansulphonic anhydride (TMSOTf) and are
 11 finally hydrolized, for example with sodium hydrogencarbonate, to give
 12 the final compound of formula (I) wherein X = O.

1 5. Process according to claim 4 wherein the possible introduction of
 2 the double bonds in position a or b is performed by reaction of
 3 dichlorodicyanoquinone (DDQ) with the corresponding silylenolethers or
 4 by oxidation with quicksilver acetate of the saturated compound
 5 obtained as claimed above and the possible transformation of the group
 6 X is performed via the corresponding enoltriflates and following
 7 carbonylation in the presence of palladium diacetate.

- 18 -

8 triphenylphosphine and the suitable nucleophilic reagent.

1 6. Compound of formula (4)



(4)

2 wherein W, Q, n, R₃, R₄, R₅ are as defined in claim 1 and Z is a
3 protecting group for the amide-group.

1 7. Pharmaceutical composition wherein the active principle is a
2 compound of formula (I) according to Claim 1 or mixtures thereof in
3 combination with the suitable pharmaceutical acceptable excipients.

1 8. Pharmaceutical composition according to Claim 7 for use in the
2 inhibition of the 5αR-I and/or 5αR-II iso-enzymes.

1 9. Pharmaceutical composition according to claims 7 and 8 in the form
2 suitable for topic use.

1 10. Use of a compound of formula (I) according to Claim 1 as a
2 medicament.

1 11. Use of a compound of formula (I) according to Claim 1 for the
2 preparation of a pharmaceutical composition for the treatment of acne,
3 baldness, prostatic cancer and prostatic hypertrophy in men and
4 hirsutism in women.

1 12. Method for the treatment of pathologies related to 5α-reductase
2 enzymes by administration to the patient of a pharmaceutically active
3 amount of a pharmaceutical composition according to Claims 7.


1 13. Method for treatment of acne, baldness, prostatic cancer and
2 prostatic hypertrophy in men and hirsutism in women, by administration

- 19 -

3 of a pharmaceutically active amount of a pharmaceutical composition
4 according to Claims 7.

1/1

SCHEME



INTERNATIONAL SEARCH REPORT

International Application No
PCT/EP 97/00552

A. CLASSIFICATION OF SUBJECT MATTER

IPC 6 C07D455/04 C07D215/22 A61K31/435

According to International Patent Classification (IPC) or to both national classification and IPC

B. FIELDS SEARCHED

Minimum documentation searched (classification system followed by classification symbols)

IPC 6 C07D

Documentation searched other than minimum documentation to the extent that such documents are included in the fields searched

Electronic data base consulted during the international search (name of data base and, where practical, search terms used)

C. DOCUMENTS CONSIDERED TO BE RELEVANT

Category *	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
X	CHEMICAL ABSTRACTS, vol. 99, no. 25, 1983 Columbus, Ohio, US; abstract no. 212524t, page 646; column 1; XP002033587 see abstract	1
X	& SU 1 027 166 A (KIEV) 7 July 1983 ---	1
X	JOURNAL OF THE CHEMICAL SOCIETY, PERKIN TRANSACTIONS 1, vol. 3, 1979, LETCHWORTH GB, pages 584-590, XP002033586 R.MORRIN ACHESON ET AL.: "ADDITION REACTIONS OF HETEROCYCLIC COMPOUNDS.PART 67." see page 584 - page 588 ---	1
-/--		

☒ Further documents are listed in the continuation of box C.

☒ Patent family members are listed in annex.

* Special categories of cited documents:

- *A* document defining the general state of the art which is not considered to be of particular relevance
- *E* earlier document but published on or after the international filing date
- *L* document which may throw doubts on priority claim(s) or which is cited to establish the publication date of another citation or other special reason (as specified)
- *O* document referring to an oral disclosure, use, exhibition or other means
- *P* document published prior to the international filing date but later than the priority date claimed

- *T* later document published after the international filing date or priority date and not in conflict with the application but cited to understand the principle or theory underlying the invention
- *X* document of particular relevance; the claimed invention cannot be considered novel or cannot be considered to involve an inventive step when the document is taken alone
- *Y* document of particular relevance; the claimed invention cannot be considered to involve an inventive step when the document is combined with one or more other such documents, such combination being obvious to a person skilled in the art.
- *&* document member of the same patent family

Date of the actual completion of the international search

23 June 1997

Date of mailing of the international search report

11.07.97

Name and mailing address of the ISA

European Patent Office, P.B. 5818 Patentlaan 2
NL - 2280 HV Rijswijk
Tel. (+31-70) 340-2040, Tx. 31 651 epo nl,
Fax: (+31-70) 340-3016

Authorized officer

Francois, J

INTERNATIONAL SEARCH REPORT

International Application No

PCT/EP 97/00552

C.(Continuation) DOCUMENTS CONSIDERED TO BE RELEVANT		
Category *	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
X	FR 1 534 278 A (BRISTOL-MYERS.) 17 June 1968 see page 1 - page 5 -----	6

INTERNATIONAL SEARCH REPORT

International application No.

PCT/EP 97/ 00552

Box I Observations where certain claims were found unsearchable (Continuation of item 1 of first sheet)

This International Search Report has not been established in respect of certain claims under Article 17(2)(a) for the following reasons:

1. ☐ Claims Nos.:
because they relate to subject matter not required to be searched by this Authority, namely:
Remark: Although claim(s) 12,13
is(are) directed to a method of treatment of the human/animal
body, the search has been carried out and based on the alleged
effects of the compound/composition.
2. ☐ Claims Nos.:
because they relate to parts of the International Application that do not comply with the prescribed requirements to such
an extent that no meaningful International Search can be carried out, specifically:
3. ☐ Claims Nos.:
because they are dependent claims and are not drafted in accordance with the second and third sentences of Rule 6.4(a).

Box II Observations where unity of invention is lacking (Continuation of item 2 of first sheet)

This International Searching Authority found multiple inventions in this international application, as follows:

1. ☐ As all required additional search fees were timely paid by the applicant, this International Search Report covers all
searchable claims.
2. ☐ As all searchable claims could be searched without effort justifying an additional fee, this Authority did not invite payment
of any additional fee.
3. ☐ As only some of the required additional search fees were timely paid by the applicant, this International Search Report
covers only those claims for which fees were paid, specifically claims Nos.:
4. ☐ No required additional search fees were timely paid by the applicant. Consequently, this International Search Report is
restricted to the invention first mentioned in the claims; it is covered by claims Nos.:

Remark on Protest

☐ The additional search fees were accompanied by the applicant's protest.

☐ No protest accompanied the payment of additional search fees.

INTERNATIONAL SEARCH REPORT

Information on patent family members

International Application No

PCT/EP 97/00552

Patent document cited in search report	Publication date	Patent family member(s)	Publication date
FR 1534278 A		BE 702549 A	12-02-68
		CH 485724 A	15-02-70
		GB 1153818 A	29-05-69
		NL 6711207 A	16-02-68
		US 3455929 A	15-07-69
